

**Case Studies in Managing Pain and Opioids:
A multimodal approach to management
(Part 1 and Part 2)**

Wendy L. Wright,
DNP, ANP-BC, FNP-BC, FAANP, FAAN, FNAP

1



Wendy L. Wright
DNP, ANP-BC, FNP-BC, FAANP, FAAN, FNAP

Adult/Family Nurse Practitioner
Owner – Wright & Associates Family Healthcare,
Amherst, NH
Owner – Partners in Healthcare Education, LLC
Faculty, Fitzgerald Health Education Associates,
Lawrence, MA

2

2

Disclosures

- Speaker Bureau:
 - Sanofi-Pasteur, Merck, Pfizer, Moderna and Seqirus: Vaccines
 - AbbVie and Biohaven: Migraines
 - Idorsia: Insomnia
 - Exact Sciences: Colorectal Cancer
 - AstraZeneca: Asthma and COPD
- Consultant:
 - Pfizer, Sanofi, Merck, Idorsia, Moderna, Seqirus, Shield Therapeutics

All relevant financial disclosures have been mitigated.

3

3

Objectives

• At the end of this presentation, the participant will be able to:

1. Describe the pathophysiology of pain as it relates to the concepts of pain management.
2. Identify evidence-based non-opioid options for the treatment of pain.
3. Discuss the risks and benefits of opioid therapy.
4. Manage ongoing opioid therapy and recognize behaviors that may be associated with opioid use disorder.

4

4

Tips



- References
 - Listed throughout and at the end of the presentation

5

5


Pathophysiology of Pain

6

6

Acute and Chronic Pain ¹		
	Acute Pain	Chronic Pain
Temporal Features	Recent, well-defined onset; expected to end in days of weeks	Remote, ill-defined onset; duration unpredictable
Biologic Function	Essential warning; impels rest and avoidance of further harm	None apparent
Intensity	Variable	Variable
Associated Affect	Anxiety common when severe or cause is unknown	Irritability or depression
Associate pain-related behaviors	Pain behaviors common when severe or cause is unknown (i.e., rubbing, splinting)	May or may not give an indication of pain
Associated features	May have signs of sympathetic hyperactivity when severe	May have vegetative signs (i.e., weight loss, loss of appetite, insomnia)
Types and examples	Post-operative trauma, burns, headache, IBS	AIDS, cancer, osteoarthritis, neuropathic pains, osteoarthritis

7



8

Physiology of Pain – Neurochemical Mediators

- A large number of neurotransmitters and other chemical mediators play a role in pain processing and modulation.
- Mediators act in a complex and interrelated process.

Excitatory (Promotes or intensifies the pain process)	Inhibitory (Blocks or dampens the pain process)
Key mediators	Key mediators
<ul style="list-style-type: none"> Glutamate Substance P Prostaglandins 	<ul style="list-style-type: none"> Opioids Norepinephrine Serotonin Gamma-aminobutyric acid (GABA)

9

Slide 7

RKO Sally, is this source (please see under slide), a "classic" or should it be updated?

Renee Kirshner, 2023-07-21T16:45:14.853

WWO 0 Please leave this - this is the citation that all use for this differentiation

Wendy Wright, 2023-07-26T12:42:24.408

RKO 1 Left as is.

Renee Kirshner, 2023-07-28T15:36:36.386

Physiology of Pain – Pathways and Effects on Pain Perception²

- Pain is a complex process mediated by multiple pathways and mechanisms both in the periphery and central nervous system (PNS and CNS [spinal cord and brain]).
- Fundamental characterization of pain
 - Nociceptive/inflammatory
 - Activation of pain-sensitive afferent neural pathways in response to injury
 - Neuropathic
 - Abnormal pain processing due to lesion of peripheral or central nervous system or both

10

10

Nociceptive Pain – Somatic²

- Somatic pain
 - Pain resulting from activation of nociceptors in the cutaneous (skin and underlying tissues) or deep tissues such as bone, blood vessels, muscles, and other supporting structures
 - **Superficial somatic pain**

Pain syndrome examples

- Traumatic bone fractures
- Muscle sprains
- Post-op incision pain

11

11

Nociceptive Pain – Visceral²

- Visceral pain
 - Activation of nociceptors in the organs and linings of the body cavities capable of responding to stimuli caused by stretching, inflammation, or ischemia to visceral structures

Pain syndrome examples

- Pancreatitis
- Hepatic metastases
- Irritable bowel syndrome

12

12

Neuropathic Pain³

- Pain believed to be sustained by aberrant somatosensory processing in the peripheral nervous system (PNS) or CNS
 - "Centrally mediated"
 - Deafferentation pain (e.g., phantom pain)
 - Sympathetically maintained pain (e.g., Complex Regional Pain Syndrome - CRPS)
 - "Peripherally mediated"
 - Originate in the nerve root, plexus, or nerve
 - Polyneuropathies and mononeuropathies

13

13

Prevalence of Neuropathic Pain




Image source: Adapted from Bennett, G.J. (1998). Neuropathic pain: new insights, new interventions. *Hosp Pract* (1995), 33(10):95-8, 101-4/4 107-10 passim. <https://pubmed.ncbi.nlm.nih.gov/9793544/>

14

14

Possible Descriptions of Neuropathic Pain

Sensations⁴

- | | |
|--|--|
| <ul style="list-style-type: none">• Numbness• Tingling• Hot-burning• Paresthetic• Paroxysmal | <ul style="list-style-type: none">• Lancing• Electric-like• Raw skin• Shooting• Deep, dull, bone-like ache |
|--|--|

15

15

Slide 14

RK0 This chart is generic enough that I don't believe it needs to be an image. The source used is from 1998. Did you want to update the info on this chart? (It's 25 yrs old of data)

Renee Kirshner, 2023-05-25T19:26:20.007

WW0 0 Renee - truly the only point of this slide is to set up the reason I speak on the top three scenarios in this lecture

Wendy Wright, 2023-06-04T18:05:08.725

RK0 1 Sally, based on what Wendy mentions above, I am going to assume that the reference at 1998 is okay and will leave as is. Let me know if that is not correct. Thanks

Renee Kirshner, 2023-07-21T15:31:21.591

WW0 2 Please leave as is -

Wendy Wright, 2023-07-26T12:42:45.746

RK0 3 Left as is.

Renee Kirshner, 2023-07-28T15:38:24.190

Slide 15

RK0 Wendy, source from 1990s. Too long ago? Alternatives to consider under the slide. Hope one works. Thanks

Renee Kirshner, 2023-07-21T16:52:15.451

WW0 0 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6003018/>

Can use this one

Wendy Wright, 2023-07-26T12:43:29.752

RK0 1 Completed.

Renee Kirshner, 2023-07-28T15:44:32.274

Possible Descriptions of Neuropathic Pain (continued)

Signs/Symptoms⁴

<ul style="list-style-type: none"> • Allodynia: Pain from a stimulus that does not normally evoke pain <ul style="list-style-type: none"> ▪ Thermal ▪ Mechanical • Hyperalgesia: Exaggerated response to a normally painful stimulus 	<ul style="list-style-type: none"> • Muscle and tissue spasm, tightness, and tenderness • Muscle weakness and atrophy • Skin color changes, rashes, swelling, and temperature abnormality
---	--

16

16

Nociceptive vs. Neuropathic Pain

Nociceptive Pain	Mixed Type	Neuropathic Pain
Caused by activity in neural pathways in response to potentially tissue-damaging stimuli ¹	Caused by a combination of both primary injury and secondary effects	Initiated or caused by primary lesion or dysfunction in the nervous system ³

17

17

TYPES OF PAIN

NOCICEPTIVE / INFLAMMATORY	NOCIPLASTIC	NEUROPATHIC	MIXED TYPES (NOCICEPTIVE / NEUROPATHIC)
Pain in response to an injury or stimuli; typically acute	Pain that arises from altered nociceptive function; typically chronic	Pain that develops when the nervous system is damaged; chronic	Primary injury and secondary effects
Post-operative pain, sports injuries, arthritis, sickle cell disease, mechanical low back pain	Fibromyalgia, irritable bowel syndrome, nonspecific low back pain	Post-herpetic neuralgia, trigeminal neuralgia, distal polyneuropathy, CRPS, neuropathic low back pain	

18

18

Slide 16

RKO Wendy, source from 1990s. Too long ago? Alternatives to consider under the slide. Hope one works. Thanks
Renee Kirshner, 2023-07-21T16:56:31.751

WWO 0 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6003018/>
Wendy Wright, 2023-07-26T12:43:36.526

RKO 1 Completed.
Renee Kirshner, 2023-07-28T15:45:07.265

BLMO
What is mixed pain?

Which of the following pain combinations are possible representations of mixed pain?

- A. Somatic and visceral pain
- B. Somatic and neuropathic pain
- C. Visceral and neuropathic pain
- D. Somatic, visceral and neuropathic pain
- E. All of the above

19

19

BLMO
What is mixed pain?

Which of the following pain combinations are possible representations of mixed pain?

- A. Somatic and visceral pain
- B. Somatic and neuropathic pain
- C. Visceral and neuropathic pain
- D. Somatic, visceral and neuropathic pain
- E. **All of the above**

20

20

Mixed Pain

- Mixed pain includes^{5,6}:
 - Specific pain syndrome such as fibromyalgia, headache syndromes, and low back pain
 - Specific disease states such as cancer or AIDS
 - Presentations of pain caused by multiple etiologies (e.g., cancer-related pain and post-herpetic neuralgia)
 - Mixed neuropathic pain is characterized by both peripherally and centrally mediated pain (e.g., stump pain from amputation and phantom limb pain).

21

21

Slide 19

DLMO Wendy- do you want this slide repeated with the answer highlighted?

Dunsmuir, Larlene M, 2023-07-24T16:03:31.998

WWO 0 UPON CLICK - ANSWER IS E - CAN HIGHLIGHT OR MAKE YELLOW

Wendy Wright, 2023-07-26T12:44:01.178

RK0 1 See next slide for answer slide.

Renee Kirshner, 2023-07-28T15:46:31.339

Slide 20

DLMO Wendy- do you want this slide repeated with the answer highlighted?

Dunsmuir, Larlene M, 2023-07-24T16:03:31.998

WWO 0 UPON CLICK - ANSWER IS E - CAN HIGHLIGHT OR MAKE YELLOW

Wendy Wright, 2023-07-26T12:44:01.178

RK0 1 completed

Renee Kirshner, 2023-07-28T15:47:17.259

What is multimodal therapy?

Which statement is true in regard to multimodal drug therapy?

1. Combines drugs and techniques that target more than one pain mechanism
2. Provides balanced and safer pain therapy
3. Incorporates into major pain management guidelines

A. Option #1
B. Option #2
C. Option #3
D. All of the above
E. None of the above

22

22

What is multimodal therapy?

Which statement is true in regard to multimodal drug therapy?

1. Combines drugs and techniques that target more than one pain mechanism
2. Provides balanced and safer pain therapy
3. Incorporates into major pain management guidelines

A. Option #1
B. Option #2
C. Option #3
D. All of the above
E. None of the above


23

23


Multimodal Therapy

24

24



25



26

Drug Therapy of Acute Pain – An Evolving Understanding

- Recognizing the need for a multimodal approach to drug therapy
 - Combinations of drugs and techniques that target more than 1 pain mechanism, not 2 drugs that target the same
 - Not a new concept, but one that is gaining increasing attention as a therapeutic framework
 - Strong evidence to support the utility of this approach; incorporated into major pain management guidelines
 - American Society of Regional Anesthesia and Pain (ASRA)
 - American Pain Society (APS)
 - American Society of Anesthesiology (ASA)

27

Slide 26

RKO Wendy, source is from 1997. (too old?) Alternatives for consideration under the slide.

Renee Kirshner, 2023-07-21T17:56:11.726


WWO 0 PLEASE LEAVE - THIS IS A PIVOTAL PAPER

Wendy Wright, 2023-07-26T12:44:24.605

RKO 1 Left as is

Renee Kirshner, 2023-07-28T15:50:15.058

POTENTIAL SITES OF ACTION FOR ANALGESIC AGENTS



28

Drug Therapy for Acute Pain – Moving to a Multimodal Strategy

- Multimodal therapy: Combining classes of agents
 - Opioids **plus** NSAIDs reduced postoperative morphine use by approximately 50%, with associated decrease in opioid-induced adverse effects and increase in patient satisfaction¹²⁻¹⁴
 - Opioid **plus** gabapentin or pregabalin reduced opioid requirements, pain, and opioid-induced adverse effects.^{15, 16}
 - Combining 2 classes of agents with opioids extends the multimodal model.

29

29

Case-based Learning
Acute Pain from End-stage OA

30

30

Clinical Case: End-stage Osteoarthritis (OA)

- Mrs. R is a 72-year-old woman presenting to her primary care provider with a 2-week history of swelling on the lateral aspect of her right knee and painless right sided foot drop.
- Imaging studies revealed end-stage osteoarthritis with swelling over the fibular head.
- The patient rated her pain a 7 out of 10.

31


Clinical Case: End-stage Osteoarthritis (OA) (continued)

- Mrs. R is a candidate for primary replacement of her knee.
 - She will undergo surgery in two weeks.
- Previously maintained on daily NSAID doses
 - Patient must discontinue NSAID in preparation for surgery to avoid interference with coagulation.

32

Osteoarthritis (OA)

- A degenerative disorder that results from breakdown of articular cartilage in the synovial joints
 - Thought to be due primarily to wear and tear
 - Non-specific inflammatory changes may also affect the joints



33

Slide 33

RKO The image is from Medscape. Thank you providing the URL. Unfortunately Medscape is copyrighted. Did you want to write for permission? If not, please change image out. Thank you. Should I use the source as a reference for the information on the slide? Please advise. Thanks,
Renee Kirshner, 2023-05-26T01:31:46.786

WWO 0 You can remove - not sure if you have any pictures of osteoarthritis knee - or joints that available = the review course has one
Wendy Wright, 2023-06-04T18:09:45.997

RKO 1 Wendy, please see far left off the slide. I found 2 possible alternative images for knees. Hope one of them works.
Renee Kirshner, 2023-07-21T15:47:09.526

WWO 2 Shutter stock image is fine - either
Wendy Wright, 2023-07-26T12:48:15.275

Clinical Case: Osteoarthritis

What is the BEST pharmacologic treatment for the patient's end-stage OA pain during the 10-day time period prior to her knee replacement surgery?

- A. Continue NSAIDs with acetaminophen at higher and more frequent doses.
- B. Opioid therapy
- C. Antidepressants
- D. Topical anesthetics
- E. None of the above

34

34

Clinical Case: Osteoarthritis

What is the BEST pharmacologic treatment for the patient's end-stage OA pain during the 10-day time period prior to her knee replacement surgery?

- A. Continue NSAIDs with acetaminophen at higher and more frequent doses.
- B. Opioid therapy
- C. Antidepressants
- D. Topical anesthetics
- E. None of the above

35

35

Pain Management Best Practices: 2019 Report²¹

- Patient-centered care in the diagnosis and treatment of acute and chronic pain
- A multimodal approach that includes medications, nerve blocks, physical therapy, and other modalities should be considered for acute pain conditions.
 - Acetaminophen can be effective for mild to moderate pain.
 - NSAIDs such as aspirin, ibuprofen, and naproxen can provide significant pain relief for inflammation, such as from arthritis, bone fractures or tumors, muscle pains, headache, and acute pain caused by injury or surgery.

36

36

Slide 35

RKO Please provide answer. Thanks
Renee Kirshner, 2023-05-26T01:38:35.489

WWO 0 There is no one correct answer - B and D are both correct but I
talk through this
Wendy Wright, 2023-06-04T18:10:28.060

RKO 1 Thanks, will leave as is.
Renee Kirshner, 2023-07-21T15:48:05.308

OTC Analgesics:

Where do they fit in pain management?

Multiple guidelines recommend OTC analgesics for the management of acute and chronic pain.

Numerous products, formulations, and delivery options are available.

Consumers can self-manage pain with OTC analgesics but are often also directed by healthcare providers to use OTC analgesics.


OTC analgesics are often more cost-effective for the consumer given the trend of higher deductibles and copays.

May be used individually or as an adjunct to RX medications

37

37

OTC Analgesics – Numerous Options



38

38

What's coming soon?

- Nerve growth factor inhibitors
 - Fasinumab
 - Tanezumab (Rejected by FDA)

39

39

**How do you want to treat
this patient?**

40

40

Case-based Learning
Postherpetic Neuralgia

41

41

Clinical Case: Cancer-related Pain and Post-herpetic Neuralgia

- Mrs. M is a 56-year-old woman with advanced breast cancer with bony metastases in the right femur and iliac crest and hepatic metastases maintained on stable doses of opioids.
- Four months ago, she developed acute herpes zoster (shingles) treated only with antiviral therapy and additional intermittent opioids with little relief.

42

42

Clinical Case: Cancer-related Pain and Post-herpetic Neuralgia (continued)

- At her medical oncologist's office, she reports steadily increasing pain in the area of her torso, unrelieved by her opioid medication.
- She states that wearing clothing over that area of her body causes excruciating pain.
 - Current pain status
 - Severe pain in the torso and upper limbs
 - Average pain intensity 6 to 8 (0 to 10)
 - Worst pain intensity 8 to 10

43

43


Clinical Case: Breast Cancer History

- Health history
 - Advanced but relatively stable breast cancer
 - Recent recovery from varicella zoster infection
- Analgesic therapy
 - Extended-release morphine 60 mg q12h
 - Short-acting morphine 15 mg q2h PRN

44

44

Schematic of VZV Latency and Reactivation



45

Slide 45

RK0 I couldn't find a source for this image on Google search. I tried accessing the URL, it went to page not found.
I did find: <https://www.micromedexsolutions.com/home/dispatch/>
(I do not have a login and didn't know if I needed to create one or if allowed)
Can you provide a different URL? Please advise. Thanks
Renee Kirshner, 2023-05-26T11:09:39.776

RK1 I did find an article that may have one or two images that may fit. However permission to use images will need to be obtained.
Please see far left for one of the images, the article and URL to gain permission. Thanks.
Renee Kirshner, 2023-05-26T11:24:58.763

WW1 0 Is there an image on zoster that is available? Picture from derm images??
Wendy Wright, 2023-06-04T18:12:21.548

RK1 1 Wendy, found the image on the slide from FNP Derm. Hope it works.
Renee Kirshner, 2023-07-21T15:55:35.664

WW1 2 fien
Wendy Wright, 2023-07-26T12:48:41.099

WW1 3 fine
Wendy Wright, 2023-07-26T12:48:44.388

Clinical Case: Breast Cancer Initial Assessment

- Patient reports excruciating pain in her torso and upper arms.
 - What type of pain is she experiencing?
 - Chronic cancer pain (somatic and visceral in origin)
 - Postherpetic neuralgia (PHN)
 - Cutaneous hypersensitivity (allodynia and hyperalgesia)
 - All of the above

46

46

Typical Locations of Herpes Zoster

- 56% thoracic
- 13% lumbar
- 13% cranial
- 11% cervical
- 4% sacral
- 3% other sites

47

47

Characterization of Pain Associated With PHN

- Dysesthesia
 - An unpleasant abnormal sensation, spontaneous or evoked^{23,24}
- Hyperalgesia
 - Pain of exaggerated severity in response to normally painful stimulation^{23,24}
- Allodynia
 - Pain evoked by a normally innocuous stimulus^{23, 24}
 - Allodynia in some patients with PHN is a form of chronic secondary hyperalgesia maintained by input from intact and possibly "irritable" primary afferent nociceptors to a sensitized CNS.²⁵

48

48

Slide 46

RK0 Is this a question that needs an answer slide or just a "talking point" to the case?

Renee Kirshner, 2023-05-26T11:28:09.976

WW0 0 It is a talking point

Wendy Wright, 2023-06-04T18:17:06.502

RK0 1 Thanks

Renee Kirshner, 2023-07-21T16:00:18.321

Slide 47

RK0 I couldn't find this image on Google search. Please provide URL. Please see far left off the slide for possible alternatives (from Shutterstock). Thanks

Renee Kirshner, 2023-05-26T11:39:26.828

WW0 0 We can use shutter stock if you keep the numbers next to it and just sub out the image

Wendy Wright, 2023-06-04T18:17:41.809

RK0 1 Let me know if there is an issue. Thanks

Renee Kirshner, 2023-07-21T16:06:03.027

WW0 2 This is great

Wendy Wright, 2023-07-26T12:49:40.354

RK0 3 Glad it works :)

Renee Kirshner, 2023-07-28T19:40:08.285

Slide 48

RK0 Wendy, 1990s source: Alternatives to consider below the slide.

Renee Kirshner, 2023-07-21T18:12:19.943

WW0 0 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5036669/#:~:text=>

<https://trialsjournal.biomedcentral.com/articles/10.1186/s13063-016>

Use both

Wendy Wright, 2023-07-26T12:49:55.095

RK0 1 completed


Renee Kirshner, 2023-07-28T16:24:42.884

PHN – Risk Factors

- Age²⁶⁻²⁸
- Severity of acute pain²⁶⁻²⁸
- Severity of acute rash²⁶⁻²⁸
- Painful prodrome²⁶
- Gender – Female²⁶

49

49



50

Clinical Case – Breast Cancer Treatment Plan

How would you manage this patient's PHN pain?

- A. Lidocaine 5% patch
- B. Opioid analgesics
- C. Tricyclic antidepressants (TCAs)
- D. Anticonvulsants
- E. Multimodal therapy

51

51

Slide 49

RKO 0 Wendy, please see source from 1990s. Alternatives to consider below the slide.

Renee Kirshner, 2023-07-21T18:18:13.059

WWO 0 Alternative sources to consider:

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4932239/>

<https://karger.com/drm/article-abstract/235/5/426/114198/Analysis>

These are fine

Wendy Wright, 2023-07-26T12:50:16.598

RKO 1 completed

Renee Kirshner, 2023-07-28T16:26:33.584

Slide 50

RKO Larlene, Please see source. It's NEJM. I can't seem to access the entire article to see if this chart was used or is actually adapted from (as stated). Normally, the chart is generic enough to let slide but it's NEJM which is a BIG NO...so I just want to verify not part of the initial article. Can you verify one way or another for me?

Thanks

Renee Kirshner, 2023-05-26T11:52:02.223

DLMO 0 Yes, I think it is generic enough and fine to use. I do wish it was more current... I have done a search and really can't find anything more current to replace it with.

Dunsmuir, Larlene M, 2023-07-24T17:50:31.955

WWO 1 Keep please

Wendy Wright, 2023-07-26T12:50:25.178

RKO 2 Left as is

Renee Kirshner, 2023-07-28T16:33:23.486

Slide 51

RKO Is an answer slide needed for this question?

Renee Kirshner, 2023-05-26T12:11:23.157

WWO 0 It is a talking point

Wendy Wright, 2023-06-04T18:18:02.888

RKO 1 Thanks

Renee Kirshner, 2023-07-21T16:08:50.519

Management Strategies for PHN²⁹

Therapy	Limitations
Lidocaine 5% patch	<ul style="list-style-type: none"> Erythema or rash Caution in patients receiving class I antiarrhythmics
Antidepressants	<ul style="list-style-type: none"> Anticholinergic AEs, sedation, cardiac conduction abnormalities
Anticonvulsants	<ul style="list-style-type: none"> Somnolence, dizziness, gait disturbances, GI upset
Opioid analgesics	<ul style="list-style-type: none"> CNS- and GI-related AEs
Dual mechanism agents	<ul style="list-style-type: none"> Similar to opioids but with better GI profile

52

Classes of Pain Medications – Local Anesthetics²²

Examples: Lidocaine, bupivacaine

- Modulate sodium channels
- When administered peripherally, may produce differential
- Also known as sensory block
 - Interrupts some nerve conduction, but leaves motor function unaffected
 - Some nerves are more readily blocked than others, depending on size and myelination.




Image source: Used with permission from Wendy Wright

53

Classes of Pain Medications – Local Anesthetics²² (continued)

Examples: Lidocaine, bupivacaine (cont.)

- Interrupts pain input at the nerve roots
- Associated with few adverse effects




Image source: Used with permission from Wendy Wright

54

Slide 52

RKO Wendy, source from 1990s. Alternatives to consider under the slide.

Renee Kirshner, 2023-07-21T18:23:00.781

WWO 0 That's because this is when the product was developed; please leave

Wendy Wright, 2023-07-26T12:50:48.378

RKO 1 Left as is

Renee Kirshner, 2023-07-28T19:40:46.527

Slide 53

RKO Could not find source for image after Google search. Please see far left off the slide for possible alternative...unless you have a URL to provide for existing image. Is this a FHEA image? Please advise. Thanks (Decision will effect slide 70, same image used)

Renee Kirshner, 2023-05-26T12:32:39.365

WWO 0 It is so generic that I don't think it an issue to use - if you have to - remove it

Wendy Wright, 2023-06-04T18:18:38.661

RKO 1 Sally, remove image and/or use "use with permission from Wendy W?"

Renee Kirshner, 2023-07-21T16:09:30.335

WWO 2 Use with permission from me

Wendy Wright, 2023-07-26T12:50:57.858

55 **Topical vs. Transdermal Medication Delivery Systems**

Topical (lidocaine patch 5%)^{3, 32-34}



Peripheral tissue activity
Applied directly over painful site
Minimal systemic absorption
Systemic AEs rare

Transdermal (fentanyl patch)^{34, 35}



Systemic activity
Applied away from painful site
Serum levels necessary
Systemic AEs common

Image sources: From British Columbia Institute of Technology (BCIT) (2017). Applying transdermal patch. https://commons.wikimedia.org/wiki/File:Applying_transdermal_patch.jpg CC BY 4.0; From Tahar, D. (2018). Generic Fentanyl Transdermal Patch. https://commons.wikimedia.org/wiki/File:A_generic_fentanyl_transdermal_patch_with_a_release_rate_of_12mcg_per_hour_applied_to_the_skin_%28cropped%29.jpg CC BY-SA 4.0. 55

55

Lidocaine Patch 5%

- Lidocaine 5% in pliable patch³⁶
- Up to 3 patches applied once daily directly over painful site^{37, 39}
 - 12 h on, 12 h off (FDA-approved label)
 - Recently published data indicate 4 patches (18–24 h) safe
- Efficacy demonstrated in 3 randomized controlled trials on PHN^{32, 38, 39}
- Drug interactions and systemic adverse effects unlikely.^{3, 32, 40}
 - Most common adverse effect: Application-site sensitivity
- Clinically insignificant serum lidocaine levels³
- Mechanical barrier decreases allodynia³⁹

56

56

Classes of Pain Medications – Antidepressants²²

- Tricyclics
 - Examples: Amitriptyline, nortriptyline, desipramine
 - Inhibit both norepinephrine (NE) and serotonin reuptake to varying degrees
 - Possesses other properties (e.g., local anesthetic-like activity)

57

57

Slide 55

RK0 Could not find sources for image during Google search. Please see far left off the slide for possible alternative; unless URLs available for original images. Thanks

Renee Kirshner, 2023-05-26T13:11:34.537

WW0 0 Yes - you can replace the images with what you have

Wendy Wright, 2023-06-04T18:19:08.045

RK0 1 Completed

Renee Kirshner, 2023-07-21T16:20:39.508

WW0 2 Just need image of patch lido and fentanyl patch

Wendy Wright, 2023-07-26T12:52:02.503

RK0 3 Images got corrupted. Please let Jill know for future. Hope this comes through now

Renee Kirshner, 2023-07-28T16:46:17.146

RK1 Wendy, please see source for #35 under the slide. Please see alternatives to consider on right under slide. (Original from 1990s)

Renee Kirshner, 2023-07-21T18:34:42.782

WW1 0 References you have found are fine -


Wendy Wright, 2023-07-26T12:51:39.116

RK1 1 completed

Renee Kirshner, 2023-07-28T19:41:26.854

Tricyclic Antidepressants – Adverse Effects²²

- Commonly reported AEs (generally anticholinergic)
 - Blurred vision
 - Cognitive changes
 - Constipation
 - Dry mouth
 - Orthostatic hypotension
 - Sedation
 - Sexual dysfunction
 - Tachycardia
 - Urinary retention




58

58

Antidepressant Use for PHN

2005 study revealed that TCAs and SSRIs reduced PHN pain, with desipramine providing satisfactory relief in 80% of those treated.


A Comparison of Pain Intensity Reduction with 3 Antidepressants



59

59

Classes of Pain Medications – Antidepressants²² (continued)



- Serotonin norepinephrine reuptake inhibitors (SNRIs)
 - Examples: Venlafaxine, duloxetine
 - Selective serotonin reuptake inhibitors (SSRIs) have not been shown to be particularly effective as pain therapy.
- Adverse effects vary by class of agent, and include dry mouth, blurred vision, nausea, constipation, agitation, dizziness, and drowsiness**

60

60

Slide 60

RKO Could not find source for image after Google search. Could not find an alternative. Is this a FHEA image? Please advise. Thanks
Renee Kirshner, 2023-05-26T13:00:06.003

WWO 0 Cite with permission from me
Wendy Wright, 2023-07-26T12:52:57.930

RKO 1 completed
Renee Kirshner, 2023-07-28T19:43:52.121

Classes of Pain Medications – Local Anesthetics²²

Examples: Gabapentin, pregabalin, lamotrigine, topiramate

- Decrease excitability of neurons by modulating sodium channels; does not act on GABA
- Emerging as top-line adjunct in acute pain and first-line therapy in chronic pain
- Adverse effects/limitations
 - Most common adverse effects are CNS related, including sleepiness, dizziness, and fatigue.




Image source: Used with permission from Wendy Wright

61

61

Treatment Plan and Outcome for Mrs. M.

- 56-year-old breast cancer patient with PHN
 - After weighing treatment options, the patient was eventually treated with multimodal therapy.
 - Continue current opioid therapy.
 - Gabapentin was given and topical lidocaine was given for local relief.

or

- Consider treatment with a single acting dual mechanism agent.
 - The patient recovered comfortably over the next 3 weeks.

62

62

Case-based Learning

Chronic Low Back Pain

63

63

Slide 61

RKO See comments from slide 62 regarding image issue

Renee Kirshner, 2023-05-26T13:26:39.780

RKO 0 Sally, delete or insert "use with permission Wendy W

Renee Kirshner, 2023-07-21T16:31:51.400

WWO 1 Permission from me

Wendy Wright, 2023-07-26T12:53:03.824

RKO 2 completed

Renee Kirshner, 2023-07-28T19:44:11.309

Clinical Case – Chronic Lower Back Pain (CLBP)

- Mr. L is 46-year-old man with history of CLBP, Type 2 diabetes, and osteoarthritis.
- Presents with an acute episode (onset 1-day prior) of low back pain
- Body mass index (BMI) – 38 kg/m²
- History of depression (currently taking sertraline)

64

64

Clinical Case – CLBP History

- Current pain status
 - Intermittent unilateral pain in the left leg with radiating weakness to the foot
 - Intensity ranges from 5/10 to 9/10
- Health history
 - Moderate osteoarthritis in the knees
 - Moderate chronic low back pain for approximately 5 years after an automobile accident

65

65

Clinical Case – CLBP History (continued)

- Medication history
 - Increasing doses of extended-release oxycodone over past year
 - Diclofenac sodium topical gel 4 g QID to each knee
 - Oxycodone extended-release 80 mg q12h with short-acting oxycodone 15 to 30 mg every 3 to 4 hours as needed

66

66

Clinical Case – CLBP Initial Assessment

Current status

- Currently patient presents with unrelieved intermittent unilateral radiating pain down the left leg and increased pain in both knees from osteoarthritis.
- Mr. L. is insisting that doses of his opioids be increased as he cannot stand the pain.
- He reports that he is tired of being on disability and wants to have a better quality of life.

67

67

Clinical Case – CLBP Initial Assessment (continued)

Identify the possible pathophysiological mechanisms for his pain.

Why is this patient **not** achieving adequate pain relief with his opioid regimen?

- A. Opioid nonresponsive neuropathic pain
- B. Opioid tolerance
- C. Worsening depression
- D. Opioid hyperalgesia
- E. Aberrant drug-seeking behaviors

RKD

68

68

Does this patient have opioid nonresponsive neuropathic pain?

Examples: Morphine, oxycodone, fentanyl

- Remains therapeutic mainstay for moderate to severe pain management²²
- Most common agents in the class act at the mu receptor.²²
- Agonistic effects both in peripheral nociceptors and centrally (spinal cord and descending pathway)²²



Image source: Used with permission from Wendy Wright

69

69

Slide 68

RK0 Please see option A. Is it "unresponsive" instead of nonresponsive?
Does this slide need an answer slide?
Renee Kirshner, 2023-05-26T14:00:42.041

WW0 0 It could be everyone o f these - no it is nonresponsive
Wendy Wright, 2023-06-04T18:19:59.306

RK0 1 Will leave as is. Thanks
Renee Kirshner, 2023-07-21T16:33:05.343

Slide 69

RK0 Could not find image on Google search. Found an alternative to consider on far left, unless you have the URL for original image or is it a FHEA image?
Renee Kirshner, 2023-05-26T14:15:06.251

WW0 0 Same as above
Wendy Wright, 2023-06-04T18:20:08.894

RK0 1 Sally, delete or use "used with permission Wendy W?
Renee Kirshner, 2023-07-21T16:33:52.968

WW0 2 Permission from me
Wendy Wright, 2023-07-26T12:54:19.362

RK0 3 completed
Renee Kirshner, 2023-07-28T19:45:01.412

Does this patient have opioid nonresponsive neuropathic pain? (continued)

Examples: Morphine, oxycodone, fentanyl (cont.)

- Considerations
 - Past hx of drug or alcohol abuse
 - Low starting dose
 - Dosing spread around the clock and not PRN



Image source: Used with permission from Wendy Wright

70

70

Is this patient developing tolerance or is pain worsening?

- Opioid tolerance is a "shift to the right" in the dose-response curve.
 - Higher dose required over time to maintain the same level of analgesia
- Tolerance can be pharmacokinetic...
 - Drug or concomitant medications upregulate metabolic pathways that remove opioids from the body




Image source: Used with permission from Wendy Wright

71

71

Is this patient developing tolerance or is pain worsening? (continued)

- ...or pharmacodynamic
 - Desensitization
 - Physiological changes to the opioid receptors
 - Downregulation
 - Internalization of opioid receptors by endocytosis, reducing their numbers




Image source: Used with permission from Wendy Wright

72

72

Slide 70

RKO Could not find image on Google search. Found an alternative to consider on far left, unless you have the URL for original image or is it a FHEA image?

Renee Kirshner, 2023-05-26T14:15:06.251

RKO 0 Sally, whatever your decision earlier, I will follow here

Renee Kirshner, 2023-07-21T16:35:16.924

WWO 1 Permission from me

Wendy Wright, 2023-07-26T12:54:46.930

Slide 71

RKO Could not find source for this image or an alternative. Can you provide an URL or is this a FHEA image? Please advise how to handle. Thanks

Renee Kirshner, 2023-05-26T14:22:18.207

RKO 0 Sally, Whatever decision made earlier, I will follow here. Thanks

Renee Kirshner, 2023-07-21T16:35:38.471

WWO 1 Permission from me or remove -

Wendy Wright, 2023-07-26T12:55:05.169

Slide 72

RKO Could not find source for this image or an alternative. Can you provide an URL or is this a FHEA image? Please advise how to handle. Thanks

Renee Kirshner, 2023-05-26T14:22:18.207

RKO 0 Sally, whatever your decision earlier, I will follow here

Renee Kirshner, 2023-07-21T16:35:58.044

WWO 1 Permission from me; these were drawn for me for a lecture I gave years ago

Wendy Wright, 2023-07-26T12:55:25.893

Is depression worsening? Psychological factors?

<ul style="list-style-type: none"> • Prolonged back pain may be associated with a psychological disturbance, manifesting as...⁴²⁻⁴⁴ <ul style="list-style-type: none"> ▪ Behavioral ▪ Cognitive ▪ Affective ▪ Somatoform (psychophysiological) 	<ul style="list-style-type: none"> • Psychological factors that may contribute to or be caused by chronic LBP include...^{42, 43} <ul style="list-style-type: none"> ▪ Depression ▪ Anxiety ▪ Somatization ▪ Posttraumatic stress disorder ▪ Pre-existing bipolar or other disorders
---	---

73

73

Social issues may contribute to chronic lower back pain (CLBP).⁴⁵

- Job dissatisfaction/loss of ability to work
- Pursuit of disability compensation
- Substance abuse
- Family dynamics
- Financial issues
- Loss of social identity or context
- Loss of ability to participate in recreational activities

74

74

Could this patient have opioid-induced hyperalgesia (OIH)?

- Increased sensitivity to pain resulting from opiate administration⁴⁶
- Opioids, in addition to providing analgesia, set in motion anti-analgesic or hyperalgesic processes⁴⁷
- Pain-free animals made tolerant to morphine have significantly decreased tolerance to pain.⁴⁷
- Opioid “tolerance” may not be a downregulation of analgesic systems, but an upregulation of hyperalgesic systems.⁴⁸

75

75

Slide 73

RK0 Did you want to write out LBP?
Renee Kirshner, 2023-05-26T14:29:35.631

RK1 Wendy, sources from 1990s. See original source on left below slide:
Please see alternatives to consider on right below slide. Thanks
Renee Kirshner, 2023-07-22T20:01:19.032

WW1 0 You can replace with alternatives below
Wendy Wright, 2023-07-26T12:55:44.927

RK1 1 completed
Renee Kirshner, 2023-07-28T19:47:27.890

Slide 74

RK0 There is a source issue here: Please see source provided on slide. I could not find any article for Wheeler and Stubbart. I could not find any article for Pathophysiology of chronic pain associated with Wheeler or Stubbart. The URL provided, redirects to: <https://emedicine.medscape.com/article/1144130-overview>, whose author is Chawla, et al and the article is low back pain and sciathica. I could use the current URL and update the author info and such or did you want to find a different article? I did find an alternative possible article at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5914385/> Let me know how you wish to handle. Thanks
Renee Kirshner, 2023-05-27T20:45:39.422

WW0 0 Alternative source to consider:
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5914385/>
Wendy Wright, 2023-07-26T12:55:56.122

RK0 1 completed
Renee Kirshner, 2023-07-28T17:33:42.285

Slide 75

RK0 Wendy, please sources 47-48. All from 1990s. Alternatives to consider under the slide.
Renee Kirshner, 2023-05-27T21:02:05.656

WW0 0 Please leave this as is
Wendy Wright, 2023-07-26T12:56:13.004

RK0 1 Left as is
Renee Kirshner, 2023-07-28T17:38:15.533

Differential Assessment⁴⁶

- General principles
 - Presence of worsening pathology or psychological influences can contribute to reports of increased pain, but are not related to opioid administration
 - Tolerance, withdrawal-related symptoms, pseudo-addiction, or addiction can be differentiated by increasing opioid dose **and/or** frequency.
 - If reports of pain increase with upward opioid titration, OIH should be considered.

76

76

Clinical Case – CLBP Initial Assessment

Identify the possible pathophysiological mechanisms for his pain.

Why is this patient **not** achieving adequate pain relief with his opioid regimen?

- A. Opioid-nonresponsive neuropathic pain
- B. Opioid tolerance
- C. Worsening depression
- D. Opioid hyperalgesia
- E. Aberrant drug-seeking behaviors

77

77

Designing an Effective Treatment Plan for Mr. L

- Initial treatment plan
 - Continue current opioid regimen (avoid escalating doses).
 - Complete opioid treatment agreement.
 - Initiate NSAID use while monitoring renal function.
 - Initiate acetaminophen use on a schedule.
 - Initiate topical analgesic use.
 - Provide patient education (body mechanics, maintaining activity).
 - Schedule physical therapy.

78

78

Slide 77

RKO Please see option A. Is it "unresponsive" instead of nonresponsive?
Does this slide need an answer slide?
Renee Kirshner, 2023-05-26T14:44:49.671

WWO 0 It is nonresponsive
Wendy Wright, 2023-06-04T18:20:27.296

RKO 1 thanks
Renee Kirshner, 2023-07-21T16:36:26.721

WAFHC Policy, per CDC 2016...

- For **all** pain patients (acute and chronic)
 - Document history and physical examination.
 - Complete opioid risk assessment tool.
 - Document treatment plan with nonpharmacologic/ pharmacologic treatments.
 - Document opioid prescription and rationale.
 - Consent form signed for opioids
 - Query the NH PDMP and print for electronic health record.

79

79

WAFHC Policy, per CDC 2016...(continued)

- **Acute** pain patients (in addition to items in I)
 - Discuss adverse effects, addiction, overdose risks.
 - Discuss risks of keeping unused medications in household.
 - Discuss options for safely securing and disposing of unused medications.
 - Discuss risks of operating heavy machinery and driving.
 - Amounts: **3 days or less**; maximum 7 days if warranted and documented rationale why 7 days is needed
 - If pain persists for more than treated time, can renew up to 30 days; however, after thirty days, must be seen for reevaluation.

80

80

WAFHC Policy, per CDC 2016... (continued)

- **Chronic** pain patients (in addition to items in I)
 - Written Treatment Agreement (Provider Patient Agreement) must be signed.
 - Refer to specialty for high risk of abuse/addiction.
 - Refer to specialty for comorbid psych disorder.
 - Query PDMP at least two times per year (ideally before every visit).
 - Random drug screening
 - In general, do not treat chronic pain in office. (Refer to subspecialty.)

81

81

2022: New CDC Guidance⁴⁹

- Removes restrictions on dosing
- Encourages providers to use best judgment
- Nonopioids first-line; but opioids when appropriate
 - Pendulum is swinging back to the middle.
- Now available
 - CDC Clinical Practice Guideline for Prescribing Opioids for Pain – United States, 2022

82

82

Initiating Opioids

- Begin with IR.
- Prescribe the lowest effective dosage.
- Use caution at any dosage, but particularly when...
 - Increasing dosage to ≥ 50 morphine milligram equivalents (MME)/day
 - Carefully justify a decision to titrate dosage to ≥ 90 MME/day.
- Always include dosing instructions, including daily maximum.
- Be aware of interindividual variability of response.
- Co-prescribe naloxone (if indicated).
- Co-prescribe bowel regimen.

83

83


Initiating Opioids (continued)

- Re-evaluate risks/benefits within 1–4 weeks (could be as soon as 3–5 days) of initiation or dose escalation
- Re-evaluate risks/benefits every 3 months; if benefits do not outweigh harms, optimize other therapies and work to taper and discontinue.

There are differences in benefit, risk and expected outcomes for patients with chronic pain and cancer pain, as well as for hospice and palliative care patients.

84

84




85

Opioid Adverse Effects

- Respiratory depression – Most serious
- Opioid-induced constipation (OIC) – Most common
- Sedation, cognitive impairment
- Falls and fractures
- Sweating, miosis, urinary retention
- Hypogonadism
- Tolerance, physical dependence, hyperalgesia
- Addiction in vulnerable patients

Prescribers should report serious AEs to the FDA:
www.fda.gov/downloads/AboutFDA/ReportsManualsForms/Forms/UCM163919.pdf or 1-800-FDA-1088



86

Opioid-induced Respiratory Depression^{50, 51}

- Chief hazard of opioid agonists, including ER/LA opioids
 - If not immediately recognized and treated, may lead to respiratory arrest and death
 - Greatest risk: Initiation of therapy or after dose increase
- Manifested by reduced urge to breathe and decreased respiration rate
 - Shallow breathing
 - CO₂ retention can exacerbate opioid sedating effects

87

Opioid

More Likely to Occur

- In elderly, cachectic, or debilitated patients
 - Contraindicated in patients with respiratory depression or conditions that increase risk
- If given concomitantly with other drugs that depress respiration
- Patients who are opioid-naïve or have just had a dose increase

Reduce Risk

- Proper dosing and titration are essential.
- Do not overestimate dose when converting dosage from another opioid product.
 - Can result in fatal overdose with first dose
- Instruct patients to swallow tablets/capsules whole.
 - Dose from cut, crushed, dissolved, or chewed tablets/capsules may be fatal, particularly in opioid-naïve individuals

88

88

WHEN TO MOVE FROM IR TO ER/LA OPIOIDS

PRIMARY REASONS

- Maintain stable blood levels (steady state plasma)
- Longer duration of action
- Multiple IR doses needed to achieve effective analgesia
- Poor analgesic efficacy despite dose titration
- Less sleep disruption

OTHER POTENTIAL REASONS

- Patient desire or need to try a new formulation
- Cost or insurance issues
- Adherence issues
- Change in clinical status requiring an opioid with different pharmacokinetics
- Problematic drug-drug interactions



89

89

Considerations for Change from IR to ER/LA Opioids⁵⁰⁻⁵³

Drug and dose selection is critical.

- Some ER/LA opioids or dosage forms are only recommended for opioid-tolerant patient.
 - Any strength of transdermal fentanyl or hydromorphone ER
 - Certain strengths/ doses of other ER/LA products (check drug prescribing information)

Monitor patients closely for respiratory depression.

- Especially within 24–72 hours of initiating therapy and increasing dosage

90

90

Considerations for Change from IR to ER/LA Opioids⁵⁰⁻⁵³
(continued)

Individualize Dosage by Titration Based on Efficacy, Tolerability, and Presence of AEs

- Check ER/LA opioid product PI for minimum titration intervals.
- Supplement with IR analgesics (opioids and non-opioid) if pain is not controlled during titration.

91

91

OPIOID TOLERANCE

If opioid tolerant, still use caution at higher doses

Patients considered opioid tolerant are taking at least

- 60 mg oral morphine/day
- 25 mcg transdermal fentanyl/hour
- 30 mg oral oxycodone/day
- 8 mg oral hydromorphone/day
- 25 mg oral oxymorphone/day
- An equianalgesic dose of another opioid



Also use caution when rotating a patient on an IR opioid to a different ER/LA opioid

Products restricted to opioid tolerant individuals include transdermal fentanyl (Duragesic) and hydromorphone (Exalgo).

SOURCE: The Opioid Analgesics Risk Evaluation & Mitigation Strategy product search, <https://opioidanalgesicsrems.com/products.html>

92

92

START WITH AN EDT FOR ADULTS



DRUG	EQUIANALGESIC DOSE		USUAL STARTING DOSE	
	SC/IV	PO	PARENTERAL	PO
Morphine	10 mg	30 mg	2.5-5 mg SC/IV q3-4hr (1.25-2.5 mg)	5-15 mg q3-4hr (IR or oral solution) (2.5-7.5 mg)
Oxycodone	NA	20 mg	NA	5-10 mg q3-4hr (2.5 mg)
Hydrocodone	NA	30 mg	NA	5 mg q3-4hr (2.5 mg)
Hydromorphone	1.5 mg	7.5 mg	0.2-0.6 mg SC/IV q2-3hr (0.2 mg)	1-2 mg q3-4hr (0.5-1 mg)


93

93

GUIDELINES FOR OPIOID ROTATION

Calculate equianalgesic dose of new opioid from EDT


REDUCE CALCULATED EQUIANALGESIC DOSE BY 25%-50%*	
SELECT % REDUCTION BASED ON CLINICAL JUDGMENT	
CLOSER TO 50% REDUCTION	CLOSER TO 25% REDUCTION
IF PATIENT... <ul style="list-style-type: none"> • Is receiving a relatively high dose of current opioid regimen • Is elderly or medically frail 	IF PATIENT... <ul style="list-style-type: none"> • Does not have these characteristics • Is changing route of administration



*75%-90% reduction for methadone

94

GUIDELINES FOR OPIOID ROTATION *(continued)*



IF SWITCHING TO METHADONE:

- Standard equianalgesic dosing tables are less helpful in opioid rotation to methadone
- For opioid tolerant patients, methadone doses should **not** exceed 30-40 mg/day upon rotation
 - Consider inpatient monitoring, including serial EKG monitoring
- For opioid-naïve patients, do **not** give methadone as an initial drug

IF SWITCHING TO TRANSDERMAL:

- **Fentanyl:** calculate dose conversion based on equianalgesic dose ratios included in the drug package insert

95

Informed Consent

Before initiating a trial of opioid analgesic therapy, confirm patient understanding of informed consent to establish:

Analgesic and Functional Goals of Treatment	The potential for and how to manage <ul style="list-style-type: none"> • Common opioid-related AEs (e.g., constipation, nausea, sedation) • Other serious risks (e.g., abuse, addiction, respiratory depression, overdose) • AEs after long-term or high-dose opioid therapy (e.g., hyperalgesia, endocrinologic or sexual dysfunction)
Expectations	
Potential Risks	
Alternatives to Opioids	

96

Consider a patient prescriber agreement (PPA).

DLM0

Reinforce expectations for appropriate and safe opioid use.

- Obtain opioids from a single prescriber.
- Fill opioid prescriptions at a designated pharmacy.
- Safeguard opioids
 - Do not store in medicine cabinet.
 - Keep locked (e.g., use a medication safe)
 - Do not share or sell medication.
- Instructions for disposal when no longer needed

97

97

Consider a patient prescriber agreement PPA. (continued)

Reinforce expectations for appropriate and safe opioid use.

- Commitments to return for follow-up visits
- Comply w/ appropriate monitoring.
 - e.g., random UDT and pill counts
- Frequency of prescriptions
- Enumerate behaviors that may lead to opioid discontinuation.
- An exit strategy

98

98

Patient Prescriber Agreement (PPA)

Document signed by both patient and prescriber at time an opioid is prescribed.

- Clarify treatment plan and goals of treatment w/patient, patient's family, and other clinicians involved in patient's care.
- Assist in patient education.
- Inform patients about the risks and benefits.
- Document patient and prescriber responsibilities.

99

99

Slide 97

DLMO ! Spelled out Patient Prescriber Agreement

Dunsmuir, Larlene M, 2023-07-24T18:28:14.670

WWO 0 fine

Wendy Wright, 2023-07-26T12:56:39.352

URINE DRUG TESTING (UDT)



- Urine testing is done **FOR** the patient, not **TO** the patient
- Helps to identify drug misuse/addiction
- Assists in assessing and documenting adherence

CLINICAL CONSIDERATIONS

- Recommend UDT before first prescription (baseline), then intermittently, depending on clinical judgment and state regulations
- Document time and date of last dose taken
- Be aware of possible false positives or negatives
- Clarify unexpected results with the lab before confronting patient to rule out poor specimen or error

100

SCREENING VERSUS CONFIRMATORY UDTs



	SCREENING (Office-based)	CONFIRMATORY (Send to lab)
Analysis technique	Immunoassay	GC-MS or HPLC
Sensitivity (power to detect a class of drugs)	Low or none when testing for semi-synthetic or synthetic opioids	High
Specificity (power to detect an individual drug)	Varies (can result in false positives or false negatives)	High
Turnaround	Rapid	Slow
Cost/Other	Lower cost; intended for a drug-free population; may not be useful in pain medicine	Higher cost; legally defensible results

GC-MS = gas chromatograph-mass spectrometry; HPLC = high-performance liquid chromatography


101

WINDOWS OF SPECIFIC DRUG DETECTION

Drug	How soon after taking drug will there be a positive drug test?	How long after taking drug will there continue to be a positive drug test?
Cannabis/ Tetrahydrocannabinol (THC)	1-3 hours	1-7 days (can be up to 1 month if long-term use)
Crack (cocaine)	2-6 hours	2-3 days
Heroin (opiates)	2-6 hours	1-3 days
Speed/upper (amphetamine, methamphetamine)	4-6 hours	2-3 days
Angel dust/PCP	4-6 hours	7-14 days
Ecstasy	2-7 hours	2-4 days
Benzodiazepine	2-7 hours	1-4 days
Barbiturates	2-4 hours	1-3 weeks
Methadone	3-8 hours	1-3 days (up to 2 weeks)
Tricyclic antidepressants	8-12 hours	2-7 days
Oxycodone	1-3 hours	1-2 days

SOURCE: <http://www.fda.gov/MedicalDevices/ProductsandMedicalProcedures/IV/IVDiagnosis/IVDiagnosis/Drugs/AboutTests/ucm125722.htm>

102



103

Management of Acute Pain⁵⁴

- "The implications for people taking opioids like morphine, oxycodone and methadone are great, since we show the short-term decision to take such opioids can have devastating consequences of making pain worse and longer lasting. This is a very ugly side to opioids that had not been recognized before."
- Study looked at just 5 days of morphine use; **increase in pain with 5 days of exposure**
- Study co-leader Prof. Linda Watkins, CU-Boulder

104

Multimodal Strategy – Implications for Practice

- Effective and safe practices with multimodal strategies require that nurses
 - Understand the rationale for combining analgesics.^{55, 56, 58}
 - Be knowledgeable about classes of analgesics.^{55, 56, 58}
 - Mechanisms of action and pharmacodynamics
 - Synergistic and AEs
 - Ensure timely administration of all analgesics, avoiding gaps in analgesia.⁵⁶⁻⁵⁸

105

Multimodal Strategy – Implications for Practice (continued)

- Effective and safe practices with multimodal strategies require that nurses (cont.)
 - Institute proper assessment and monitoring practices^{56, 57}
 - Aggressively manage AEs of analgesics^{55, 56, 58}
 - Remain informed about novel dual-mechanism analgesics and drug delivery systems^{55, 56, 58}

106

A CLOSER LOOK AT THE ORT-ODU

Opioid Risk Tool – OUD (ORT-ODU)

This tool should be administered to patients upon an initial visit prior to beginning or continuing opioid therapy for pain management. A score of 2 or lower indicates low risk for future opioid use disorder; a score of >= 3 indicates high risk for opioid use disorder.

Mark X with Yes (or apply)	YES	NO
Family history of substance abuse		
Alcohol	1	0
Illegal drugs	1	0
Rx drugs	1	0
Personal history of substance abuse		
Alcohol	1	0
Illegal drugs	1	0
Rx drugs	1	0
Has history of 3 or more	1	0
Psychological distress		
ADDC OUDS, bipolar, schizophrenia	1	0
Depression	1	0
Smoking index	1	0

Substance use disorder history does not prohibit treatment with opioids but may require additional monitoring and expert consultation or referral.

Scoring:

- ≤ 2: low risk
- ≥ 3: high risk

SOURCE: Cheetham, M., et al. J Pain 2019, Jan 26

107

Naloxone⁵⁹

- Every person using opioids should have naloxone available.
- Now sold over the counter without a prescription
- Illicit opioids may be associated with rigid chest wall syndrome requiring multiple doses.

108

When should patients be referred to a pain management specialist?⁴⁹

<ul style="list-style-type: none"> • Complex pain syndromes • Unsuccessful outcomes • Multimodal therapy • History or pre-existing substance abuse 	<ul style="list-style-type: none"> • Problems with adherence • Interventional procedures • Behavioral or cognitive therapy
--	---

109

109

Drug-related Behaviors that Need to be Evaluated⁶⁰

Probably Less Predictive	Probably More Predictive
<ul style="list-style-type: none"> • Aggressive complaining • Medication hoarding when symptoms milder • Requesting specific medications • Acquisition of medications from other medical sources • Unsanctioned dose escalation once or twice 	<ul style="list-style-type: none"> • Selling prescription medications • Prescription forgery • Stealing or "borrowing" medications from another person • Injecting oral formulation • Obtaining prescription medications from nonmedical source • Multiple episodes of prescription "loss"

110

110

Drug-related Behaviors that Need to be Evaluated⁶⁰ (continued)

Probably Less Predictive	Probably More Predictive
<ul style="list-style-type: none"> • Unapproved use of the medication to treat another symptom • Reporting psychic effects not intended by the clinician • Occasional impairment 	<ul style="list-style-type: none"> • Concurrent abuse of related illicit drugs • Multiple dose escalations despite warnings • Repeated episodes of gross impairment or dishevelment

111

111

Slide 110

RKO Wendy, please see source on slide 1990s. Alternatives to consider under the slide.

Renee Kirshner, 2023-07-21T19:15:18.557

WWO 0 <https://nida.nih.gov/research-topics/parents-educators/conversati>

Wendy Wright, 2023-07-26T12:56:57.200

RKO 1 completed

Renee Kirshner, 2023-07-28T19:50:35.773

I would be happy to entertain
any questions or comments

112

112

Thank you for your time and attention.

Wendy L. Wright,
DNP, ANP-BC, FNP-BC, FAANP, FAAN, FNAP

WendyARNP@aol.com

113

113

^{KS}
References

1. Portenoy, R.K., Kanner, R.M. (1996). Definition and Assessment of Pain. In: Portenoy, R.K., Kanner, R.M., eds. *Pain Management: Theory and Practice*. Philadelphia, Pa: FA Davis Company.
2. Galer, B., Gammaitoni, A., Alvarez, N. (2003). Pain. In: Dale DC, Federman DD, eds. *WebMD Scientific American® Medicine*. New York, NY:WebMD Corporation.
3. Galer, B.S., Dworkin, R.H. (2000). *A Clinical Guide to Neuropathic Pain*. New York, NY: McGraw Hill, Healthcare Information Programs.
4. Murnion, B.P. (2018). Neuropathic pain: current definition and review of drug treatment. *Aust Prescr.*, 41(3):60-63. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6003018/>
5. Portenoy, R.K., Bennett, D.S., Rauck, R., Simon, S., Taylor, D., Brennan, M., Shoemaker, S. (2006). Prevalence and characteristics of breakthrough pain in opioid-treated patients with chronic noncancer pain. *J Pain*, 7(8):583-91. <https://pubmed.ncbi.nlm.nih.gov/16885015/>
6. Webster, L.R. (2008). Breakthrough pain in the management of chronic persistent pain syndromes. *Am J Manag Care*, 14(5 Suppl 1):S116-22. <https://pubmed.ncbi.nlm.nih.gov/18611099/>

114

114

Slide 114

RKO Sally, many of these sources are from mid 90. What is the cut off for updated sources?

Renee Kirshner, 2023-05-26T20:30:27.141

WWO 0 See all of the changes in the program

Wendy Wright, 2023-07-26T12:57:12.135

RKO 1 Left many sources in 1990s as indicated.

Renee Kirshner, 2023-07-28T19:51:11.854

References (continued)

7. National Pharmaceutical Council Inc. (2001). *Pain: Current Understandin of Assessment, Management, and Treatments*. <https://www.npcnow.org/sites/default/files/media/Pain-Current-Understanding-of-Assessment-Management-and-Treatments.pdf>
8. Gordon, D.B., Dahl, J.L., Miaskowski, C., McCarberg, B., Todd, K.H., Paice, J.A., Lipman, A.G., Bookbinder, M., Sanders, S.H., Turk, D.C., Carr, D.B. (2005). American pain society recommendations for improving the quality of acute and cancer pain management: American Pain Society Quality of Care Task Force. *Arch Intern Med.*, 165(14):1574-80. <https://pubmed.ncbi.nlm.nih.gov/16043674/>
9. Authors unknown. (1997). Practice guidelines for chronic pain management. A report by the American Society of Anesthesiologists Task Force on Pain Management, Chronic Pain Section. *Anesthesiology*, 86(4):995-1004. <https://pubmed.ncbi.nlm.nih.gov/9105246/>
10. Rathmell, J.P., Wu, C.L., Sinatra, R.S., Ballantyne, J.C., Ginsberg, B., Gordon, D.B., Liu, S.S., Perkins, F.M., Reuben, S.S., Rosenquist, R.W., Viscusi, E.R. (2006). Acute post-surgical pain management: a critical appraisal of current practice, December 2-4, 2005. *Reg Anesth Pain Med.*, 31(4 Suppl 1):1-42. <https://pubmed.ncbi.nlm.nih.gov/16849098/>

115

115

References (continued)

11. American Society of Anesthesiologists Task Force on Acute Pain Management. (2004). Practice Guidelines for Acute Pain Management in the Perioperative Setting: An Updated Report by the American Society of Anesthesiologists Task Force on Acute Pain Management. *Anesthesiology*, 100:1573-1581. <https://pubs.asahq.org/anesthesiology/article/100/6/1573/6558/Practice-Guidelines-for-Acute-Pain-Management-In>
12. Stephens, J., Laskin, B., Pashos, C., Peña, B., Wong, J. (2003). The burden of acute postoperative pain and the potential role of the COX-2-specific inhibitors. *Rheumatology* (Oxford), 42 Suppl 3:iii40-52. <https://pubmed.ncbi.nlm.nih.gov/14585917/>
13. Zimmel, M.H. (2006). The role of COX-2 inhibitors in the perioperative setting: efficacy and safety. A systematic review. In: Database of Abstracts of Reviews of Effects (DARE): Quality-assessed Reviews [Internet]. York (UK): Centre for Reviews and Dissemination (UK) <https://www.ncbi.nlm.nih.gov/books/NBK73241/>
14. Tiippana, E.M., Hamunen, K., Kontinen, V.K., Kalso, E. (2007). Do surgical patients benefit from perioperative gabapentin/pregabalin? A systematic review of efficacy and safety. *Anesth Analg.*, 104(6):1545-56, table of contents. <https://pubmed.ncbi.nlm.nih.gov/17513656/>

116

116

References (continued)

15. Mathiesen, O., Moinicke, S., Dahl, J.B. (2007). Gabapentin and postoperative pain: a qualitative and quantitative systematic review, with focus on procedure. *BMC Anesthesiol.*, 7:6. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1950698/>
16. Elia, N., Lysakowski, C., Tramèr, M.R. (2005). Does multimodal analgesia with acetaminophen, nonsteroidal antiinflammatory drugs, or selective cyclooxygenase-2 inhibitors and patient-controlled analgesia morphine offer advantages over morphine alone? Meta-analyses of randomized trials. *Anesthesiology*, 103(6):1296-304. <https://pubmed.ncbi.nlm.nih.gov/16306743/>
17. Tzschentke, T.M., De Vry, J., Terlinden, R. (2006). Tapentadol hydrochloride. Analgesic, mu-opioid receptor agonist, noradrenaline reuptake inhibitor. *Drugs Future*, 31:1053-1061
18. FDA. (2010). Tapentadol (Nucynta®). Prescribing information. https://www.accessdata.fda.gov/drugsatfda_docs/label/2010/022304s003lbl.pdf.
19. Kleiner, R. & Hallmann, Caspar & Steup, Achim. (2006). Efficacy of a single dose of tapentadol HCl for analgesia after third molar surgery. *J Pain*, 7. S44. Abstract 773

117

117

References (continued)

20. American Society of Anesthesiologists Task Force on Chronic Pain Management; American Society of Regional Anesthesia and Pain Medicine. (2010). Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology*, 112(4):810-33. <https://pubs.asahq.org/anesthesiology/article/112/4/810/10691/Practice-Guidelines-for-Chronic-Pain-ManagementAn>
21. U.S. Department of Health and Human Services. (2019). Pain Management Best Practices Inter-Agency Task Force Report: Updates, Gaps, Inconsistencies, and Recommendations. <https://www.hhs.gov/sites/default/files/pmtf-final-report-2019-05-23.pdf>
22. Brunto, L.L., Knollman, B.C., Hilal-Dandan, R. (2017). *Goodman & Gilman's The Pharmacological Basis of Therapeutics*, (13th ed.). McGraw Hill Medical.
23. Mallick-Searle T, Snodgrass B, Brant JM. (2016). Postherpetic neuralgia: epidemiology, pathophysiology, and pain management pharmacology. *J Multidiscip Healthc.*, 9:447-454. [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5036669/#:~:text=Postherpetic%20neuralgia%20\(PHN\)%20is%20a,herpes%20zoster%20\(HZ\)%20rash](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5036669/#:~:text=Postherpetic%20neuralgia%20(PHN)%20is%20a,herpes%20zoster%20(HZ)%20rash).

118

118

References (continued)

24. Rullán, M., Bullete, O., Leiva, A., Soler, A., Roca, A., Gonzalez-Bals, M.J., Lorente, P., Llobera, J. and PHN group. (2017). Efficacy of gabapentin for prevention of postherpetic neuralgia: study protocol for a randomized controlled clinical trial. *Trials* 18, 24. <https://trialsjournal.biomedcentral.com/articles/10.1186/s13063-016-1729-y>
25. Petersen, K. L., Fields, H. L., Brennum, J., Sandroni, P., & Rowbotham, M. C. (2000). Capsaicin evoked pain and allodynia in post-herpetic neuralgia. *Pain*, 88(2), 125-133.
26. Jung, B. F., Johnson, R. W., Griffin, D. R., & Dworkin, R. H. (2004). Risk factors for postherpetic neuralgia in patients with herpes zoster. *Neurology*, 62(9), 1545-1551.
27. Forbes, H.J., Bhaskaran, K., Thomas, S.L., Smeeth, L., Clayton, T., Mansfield, K., Minassian, C., Langan, S.M. (2016). Quantification of risk factors for postherpetic neuralgia in herpes zoster patients: A cohort study. *Neurology*, 87(1):94-102. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4932239/>
28. Wei, S., Li, X., Wang, H., Liu, Q., Shao, L. (2019). Analysis of the Risk Factors for Postherpetic Neuralgia. *Dermatology*, 235 (5): 426-433. <https://karger.com/drm/article-abstract/235/5/426/114198/Analysis-of-the-Risk-Factors-for-Postherpetic?redirectedFrom=fulltext>

119

119

References (continued)

29. Kost, R.G., Strauss, S.E. (1996). Postherpetic neuralgia—pathogenesis, treatment, and prevention. *N Engl J Med.*, 335:32-42.
30. Oster, G., Harding, G., Dukes, E., Edelsberg, J., & Cleary, P. D. (2005). Pain, medication use, and health-related quality of life in older persons with postherpetic neuralgia: results from a population-based survey. *The Journal of pain*, 6(6), 356-363.
31. Oxman, M. N., Levin, M. J., Johnson, G. R., Schmader, K. E., Straus, S. E., Gelb, L. D., Arbeit, R.D., Simberkoff, M.S., Gershon, A.A., Davis, L.E., Weinberg, A., Boardman, K.D., Williams, H.M., Zhang, J.H., Peduzzi, P.N., Beisel, C.E., Morrison, V.A., Guatelli, J.C.,... & Silber, J. L.; Shingles Prevention Study Group. (2005). A vaccine to prevent herpes zoster and postherpetic neuralgia in older adults. *New England Journal of Medicine*, 352(22), 2271-2284.
32. Argoff, C.E. (2000). New analgesics for neuropathic pain: the lidocaine patch. *Clin J Pain*, 16(2 Suppl):S62-6. <https://pubmed.ncbi.nlm.nih.gov/10870742/>
33. Gudin, J. and Nalamachu, S. (2020) Utility of lidocaine as a topical analgesic and improvements in patch delivery systems. *Postgraduate Medicine*, 132:1. 28-36, <https://www.tandfonline.com/doi/full/10.1080/00325481.2019.1702296?scroll=top&needAccess=true&ole=tab>

120

120

References (continued)

- 34. Burch, F., Coddling, C., Patel, N., Sheldon, E. (2004). Lidocaine patch 5% improves pain, stiffness, and physical function in osteoarthritis pain patients. A prospective, multicenter, open-label effectiveness trial. *Osteoarthritis Cartilage*, 12(3):253-5. [https://www.oarsjournal.com/article/S1063-4584\(03\)00272-3/fulltext](https://www.oarsjournal.com/article/S1063-4584(03)00272-3/fulltext)
- 35. Fentanyl transdermal system (Duragesic®). (2021). Prescribing information. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/019813s081lbl.pdf
- 36. FDA. (2015). Lidocaine patch 5% (Licoderm®). Prescribing information. https://www.accessdata.fda.gov/drugsatfda_docs/label/2015/020612s012lbl.pdf. Endo Pharmaceuticals Inc.
- 37. Gammaitoni, A.R., Alvarez, N.A., Galer, B.S. (2002). Pharmacokinetics and safety of continuously applied lidocaine patches 5%. *Am J Health Syst Pharm.*, 59(22):2215-20. <https://pubmed.ncbi.nlm.nih.gov/12455305/>

121

References (continued)

- 38. Cheville, A.L., Sloan, J.A., Northfelt, D.W., Jillella, A.P., Wong, G.Y., Bearden Iii, J.D., Liu, H., Schaefer, P.L., Marchello, B.T., Christensen, B.J., Loprinzi, C.L. (2009). Use of a lidocaine patch in the management of postsurgical neuropathic pain in patients with cancer: a phase III double-blind crossover study (N01CB). *Support Care Cancer*, 17(4):451-60. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5653964/>
- 39. Hines, R., Keaney, D., Moskowitz, M.H., Prakken, S. (2002). Use of Lidocaine Patch 5% for Chronic Low Back Pain: A Report of Four Cases. *Pain Medicine*. Volume 3, Issue 4, Pages 361–365. <https://academic.oup.com/painmedicine/article/3/4/361/11843399>
- 40. Gammaitoni, A. R., & Davis, M. W. (2002). Pharmacokinetics and tolerability of lidocaine patch 5% with extended dosing. *Annals of Pharmacotherapy*, 36(2), 236-240. <https://pubmed.ncbi.nlm.nih.gov/11847940/>
- 41. Kalso, E., Allan, L., Dobrogowski, J., Johnson, M., Krceviski-Skvarc, N., Macfarlane, G. J., Mick, S., Ortolani, S., Perrot, S., Perucho, A., Semmons, I. and Sørensen, J. (2005). Do strong opioids have a role in the early management of back pain? Recommendations from a European expert panel. *Current medical research and opinion*, 21(11), 1819-1828

122

References (continued)

- 42. American Psychological Association. (2019) *APA Clinical Practice Guideline for the Treatment of Depression Across Three Age Cohorts*. <https://www.apa.org/depression-guideline/guideline.pdf>
- 43. Singhal, K., Muliya, K.P., Pakhare, A.P., Behera, P., Santoshi, J.A. (2021). Do Patients of Chronic Low Back Pain have Psychological Comorbidities? *Avicenna J Med.*, 11(3):145-151. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8500070/>
- 44. Robertson, D., Kumbhare, D., Nolet, P., Srbely, J., Newton, G. (2017). Associations between low back pain and depression and somatization in a Canadian emerging adult population. *J Can Chiropr Assoc.*, 61(2):96-105. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5596967/>
- 45. Makris, U.E., Higashi, R.T., Marks, E.G., Fraenkel, L., Gill, T.M., Friedly, J.L., Reid, M.C. (2017). Physical, Emotional, and Social Impacts of Restricting Back Pain in Older Adults: A Qualitative Study. *Pain Med.*, 18(7):1225-1235. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5914385/>
- 46. Tompkins, D.A., Campbell, C.M. (2011). Opioid-induced hyperalgesia: clinically relevant or extraneous research phenomenon? *Curr Pain Headache Rep.*, 15(2):129-36. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3165032/>

123

References (continued)

- 47. Compton, M.A. (1994). Cold-pressor pain tolerance in opiate and cocaine abusers: correlates of drug type and use status. *J Pain Symptom Manage.*, 9(7):462-73. <https://pubmed.ncbi.nlm.nih.gov/7822886/>
- 48. Laulin, J.P., Célérier, E., Larcher, A., Le Moal, M., Simonnet, G. (1999). Opiate tolerance to daily heroin administration: an apparent phenomenon associated with enhanced pain sensitivity. *Neuroscience*, 89(3):631-6. <https://pubmed.ncbi.nlm.nih.gov/10199599/>
- 49. CDC. (2022). CDC Clinical Practice Guideline for Prescribing Opioids for Pain — United States, 2022 *MMWR*, 71(3): 1–95. <https://www.cdc.gov/mmwr/volumes/71/rr/rr7103a1.htm>
- 50. Chou, R., Fanciullo, G.J., Fine, P.G., Adler, J.A., Ballantyne, J.C., Davies, P., Donovan, M.I., Fishbain, D.A., Foley, K.M., Fudin, J., Gilson, A.M., Ketter, A., Mauskop, A., O'Connor, P.G., Passik, S.D., Pasternak, G.W., Portenoy, R.K., Rich, B.A., Roberts, R.G., Todd, K.H., Miaskowski, C.; American Pain Society-American Academy of Pain Medicine Opioids Guidelines Panel. (2009). Clinical guidelines for the use of chronic opioid therapy in chronic noncancer pain. *J Pain.*, 10(2):113-30. <https://pubmed.ncbi.nlm.nih.gov/19187889/>

124

124

References (continued)

- 51. FDA. (2018). *Education Blueprint for Health Care Providers Involved in the Treatment and Monitoring of Patients with Pain*. <https://www.fda.gov/media/99496/download>
- 52. FDA. (2017). *Introduction for the FDA Blueprint for Prescriber Education for Extended-Release and Long-Acting Opioid Analgesics*. <https://www.fda.org/2017/Blueprint%20Opioid%20LA.ER%20REMS%20as%20of%201.20.2017.pdf>
- 53. Whiteman, H. (2016) *Opioids might worsen chronic pain, study finds*. Medical News Today website. <https://www.medicalnewstoday.com/articles/310645>
- 54. Krenzischek, D.A., Dunwoody, C.J., Polomano, R.C., Rathmell, J.P. (2008). Pharmacotherapy for acute pain: implications for practice. *Pain Manag Nurs.*, 9(1 Suppl):S22-32. <https://pubmed.ncbi.nlm.nih.gov/18294591/>
- 55. Dunwoody, C. J., Krenzischek, D. A., Pasero, C., Rathmell, J. P., & Polomano, R. C. (2008). Assessment, physiological monitoring, and consequences of inadequately treated acute pain. *Pain Management Nursing*, 9(1), 11-21.
- 56. Polomano, R.C., Dunwoody, C.J., Krenzischek, D.A., Rathmell, J.P. (2008). Perspective on pain management in the 21st century. *Pain Manag Nurs.*, 9(1 Suppl):S3-10. <https://pubmed.ncbi.nlm.nih.gov/18294589/>

125

125

References (continued)

- 57. Polomano, R.C., Rathmell, J.P., Krenzischek, D.A., Dunwoody, C.J. (2008). Emerging trends and new approaches to acute pain management. *Pain Manag Nurs.*, 9(1 Suppl):S33-41. <https://pubmed.ncbi.nlm.nih.gov/18294592/>
- 58. Cheattle, M.D., Compton, P.A., Dhingra, L., Wasser, T.E., O'Brien, C.P. (2019). Development of the Revised Opioid Risk Tool to Predict Opioid Use Disorder in Patients with Chronic Nonmalignant Pain. *J Pain*, 20(7):842-851. <https://pubmed.ncbi.nlm.nih.gov/30690168/>
- 59. CDC. (2023). *When to Offer Naloxone to Patients*. https://www.cdc.gov/opioids/naloxone/factsheets/pdf/Naloxone_FactSheet_Clinicians.pdf
- 60. NIH on Drug Abuse (2021). *What Are the Signs of Having a Problem With Drugs?* <https://nida.nih.gov/research-topics/parents-educators/conversation-starters/what-are-signs-having-problem-drugs>

CO*RE slides used with permission, obtained by Wendy Wright.
 Slides include: 8, 18, 25, 28, 96, 101, 104-107, 112-115, 119

Unless noted, images are from Shutterstock, used with permission from Colibri Group Healthcare license.

126

126